

**RESISTANCE STATUS OF Aedes Aegypti TO INSECTICIDES IN THE JAZAN REGION OF SAUDI ARABIA****A. A. ALSHEIKH, W. S. MOHAMMED, \*E. M. NOURELDIN, O. M. DAFFALLA, K. J. SHRWANI, Y. A. HOBANI, F. A. ALSHEIKH<sup>1</sup>, M. H. ALZHRANI<sup>2</sup> AND A. A. BINSAEED<sup>3</sup>**National Center for vector-borne Diseases,  
<sup>1</sup>College of Public Health and Tropical Medicine,  
Jazan University. SAUDI ARABIA.<sup>2</sup>Malaria program Directorate, SAUDI ARABIA<sup>3</sup>Ministry of Public Health,  
MoH- RIYADH, SAUDI ARABIA

\*Corresponding Author

Email : siddignoureldin@hotmail.com

**Received : 12.2.16; Accepted : 28.3.16****ABSTRACT**

The present study was designed to provide information on the susceptibility status of the adult and larvae of *Aedes aegypti* mosquitoes in Jazan region of Saudi Arabia. Bioassay tests were performed on adults and larvae by using WHO recommended concentrations and test kits. Adults of *A. aegypti* mosquitoes were exposed to test papers impregnated with Lambda-cyhalothrin (0.05%), Cyfluthrin (0.15%), Deltamethrin (0.05%), Permethrin (0.75%), Fenitrothion (1%), Bendiocarb (0.1%) and DDT (4%) insecticides. *A. aegypti* mosquitoes were found to be susceptible only to Cyfluthrin; (mortality rate was 100 %), whereas variable resistances were observed from the rest of the other insecticides tested (mortality rates ranged between 93.6 and 17%). Larvae were subjected to different concentrations of Diflubenzuron, Methoprene (IGRs) and Temephos (Organophosphate). The inhibition of adult emergence ( $IE_{50}$  and  $IE_{95}$ ) values for the IGRs and the ( $LC_{50}$  and  $LC_{95}$ ) for Temephos have been determined by log-probit regression analysis. *A. aegypti* larvae were resistant to Temephos ( $LC_{50}$  61.8 -  $LC_{95}$  35600.1 mg/l) and showed high susceptibility to Methoprene than Diflubenzuron ( $IE_{50}$  0.49 -  $IE_{95}$  10.9 mg/l) and ( $IE_{50}$  0.86 and  $IE_{95}$  93.8 mg/l), respectively. The larvae were more susceptible to Methoprene than Diflubenzuron by 1.8 folds.

Figure : 00

References : 41

Tables : 03

KEY WORDS : *Aedes aegypti*, Insecticides, Jazan region, Resistance, Saudi Arabia.**Introduction**

Dengue virus (DENV, Flaviviridae, Flavivirus) and Chikungunya virus (CHIKV, Togaviridae, Alphavirus) are the most significant human viral pathogens spread by the bite of an infected mosquito in most tropical regions. The annual reported cases are estimated to be 50-100 million, including 500,000 severe cases of dengue hemorrhagic fever (DHF) or dengue shock syndrome (DSS)<sup>35</sup>. DENV are the most prevalent mosquito-borne human virus worldwide.<sup>11</sup> Globally, *Aedes aegypti* and *Aedes albopictus* are the main epidemic vectors of DENV and CHIKV.<sup>14,24</sup>

No vaccine or antiviral therapy currently available, disease prevention relies largely on surveillance and mosquito control by using

insecticides.

Pyrethroids are still the mainstay of vector control program. The use of pyrethroid insecticides has, however, increased and that of the organochlorines and some of the more toxic organophosphate compounds have decreased in recent years.

The extensive use of insecticides in agriculture, house spraying and space spraying result in the appearing of insect that can survive and make resistance to insecticides. Appropriate monitoring of vector resistance to insecticides is an integral part of planning and evaluation of insecticide uses in control programs.

Unfortunately, many vector control programs are threatened by the development of insecticide

resistance in *A. aegypti* and *A. albopictus*.<sup>21,41</sup> Resistance to multiple insecticides (e.g. pyrethroids and organophosphates) has been reported in *A. aegypti* in South America and the Caribbean.<sup>21,25</sup> DDT resistance is now widely distributed in *A. aegypti* throughout northern Thailand and it is also resistant to the pyrethroids, permethrin and deltamethrin, in many areas.<sup>26,32</sup>

In India, most of the studies reviewed have revealed that the adult *Aedes aegypti* and *Aedes albopictus* mosquitoes are resistant to DDT, but generally, remain susceptible to malathion, temephos, propoxur and fenitrothion. The larval stages of *Aedes aegypti* and *Aedes albopictus* have also been found to be resistant to DDT, but susceptible to larvicides, namely temephos, fenthion, and malathion.<sup>31</sup>

In South India, workers<sup>29</sup> reported that adult *A. aegypti* and *A. albopictus* were resistant to DDT and dieldrin, but susceptible to propoxur, fenitrothion, malathion, deltamethrin, permethrin and lambda-cyhalothrin. The resistance to pyrethroids (permethrin and deltamethrin) has been developed in *A. aegypti* in most of the collected areas of Thailand.<sup>13</sup> Workers have tested *Aedes aegypti*, which is the main dengue vector in Baan Suan community (Thailand) with deltamethrin, permethrin and fenitrothion.<sup>26</sup> Their results showed that dengue mosquitoes are clearly resistant to permethrin and tolerant to deltamethrin, but were 100% susceptible to fenitrothion.

In Costa Rica, the *A. aegypti* larvae showed higher resistance levels to the organophosphate (OP) temephos and the adult were resistant to deltamethrin, but susceptible to bendiocarb, chlorpyrifos, and cypermethrin.<sup>5</sup> In Havana city, a high resistance of *A. aegypti* larvae to temephos and fenthion was observed.<sup>20</sup>

Currently, *A. aegypti* resistance to temephos is widespread in the Americas.<sup>36</sup> Acquired resistance by *A. aegypti* to temephos was reported

in Brazil<sup>6,17,19</sup>, Thailand<sup>12</sup>, Bolivia and Argentina.<sup>4</sup> Recently, incipient resistance to temephos was also reported in *A. aegypti* in Argentina.<sup>27</sup>

In Malaysia, *Aedes* larvae tolerance against temephos has been reported and have exhibited moderate resistance toward methoprene and low resistance toward pyriproxyfen<sup>9,16</sup>, but susceptible to diflubenzuron, cyromazine, and novaluron.<sup>15</sup>

During the past two decades, considerable progress has been made in the development of natural and synthetic compounds known as insect growth regulators (IGRs) which are capable of interfering with the process of growth, development, and metamorphosis of the target mosquito species. Two types of IGRs are available, one which inhibits the growth of larvae (Methoprene and Pyriproxyfen) due to juvenile hormone-like action and known as JH mimics or analogs and the other type of IGR compound (Diflubenzuron) which interferes with chitin production leading to moulting disturbances, resulting in the death of the insect.

In Jeddah, Saudi Arabia, the LC<sub>50</sub> of five insecticides against *A.aegypti* larvae have been evaluated.<sup>2</sup> The results revealed that the larvae proved to be more susceptible to Lambda-cyhalothrin (0.01 ppm) than Snap >γPermethrin 11% + Tetramethrin 1% + Piperonyl butoxide 11% >γ (0.048 ppm) and Bacilod >γ*Bacillus thuringiensis israelensis*?γ (0.3 ppm) by about 4.8 and 30 folds, respectively. Moreover, Baycidal >γTriflumuron 25WP?γ (0.0007 ppm) proved to be more effective against *A. aegypti* than Sumilarv >γPyriproxyfen 0.5g ?γ(0.003 ppm) by about 4.3 folds.

In Makkah city, Saudi Arabia, it was found that the Lambdacyhalothrin was the most effective larvicide against *A. aegypti* with LC<sub>50</sub> values of 0.007 ppm and 0.012ppm for the laboratory and field strains, respectively.<sup>3</sup>

In Jazan region of Saudi Arabia, there are no available data on pesticide resistance for *A. aegypti* against IGR and chemical insecticides

**TABLE- 1: Efficacy of Temephos against *A. aegypti* larvae**

Insecticide	Concentration (mg/l)	Larval Mortality (%)	LC <sub>50</sub> (LC <sub>95</sub> ) (mg/l)	95% Confidence limit	
				Lower LC <sub>50</sub> (LC <sub>95</sub> )	Upper LC <sub>50</sub> (LC <sub>95</sub> )
Temephos	0.005 – 0.625	0 - 10	61.8 (35600.1)	3.2 (101.9)	8.2 (1.31E+46)

used in the area. In order to implement effective and sustainable arbovirus vector control measures, there is an urgent need to determine the susceptibility of the major vector of DENV and CHIK to insecticides commonly used for mosquito control. We, therefore, examined the insecticide susceptibility for the adult and Larvae of *A. aegypti* populations in Jazan region using WHOES-approved procedures.<sup>37,38</sup>

### Materials and Methods

This study was carried out in Jazan region which is located in Southwest Saudi Arabia, lies between 16°-12, and 18°-25, N latitude. The Jazan region is situated in the subtropical zone and has an average monthly temperature ranging between 25.8°C in January to 33.4°C in July and average relative humidity ranges between 55% and 72.5%, with an average rain of 77 - 56.7 mm, (August and October), respectively.<sup>29</sup>

#### Mosquitoes

*Aedes aegypti* larvae were collected from different parts of Jazan region and brought to the insectory of the National Center for Vector-Borne Disease in Gizan city. The larvae were kept under control conditions of temperature ( $25 \pm 2$ ) °C and relative humidity (70-80%) with a photo period of 12-hour darkness and 12-hour light. Pupae from water medium were transferred to mosquito rearing cages (30 cm<sup>3</sup>). Resulting mosquito adults were provided with 10% glucose solution supplemented with 1% vitamin B complex soaked in cotton wool.

Female mosquitoes were permitted to blood-feed on pigeons after 5 days of adult emergence. After 3 days of feeding, a piece of moist filter paper in a porcelain bowl half filled with water was introduced for egg deposition.

Eggs from filial generation ( $F_1$ ) were hatched and larvae were reared in plastic trays and fed every alternate day with a powdered mixture of wheat, yeast and milk (1:1:1). Late third or early fourth instar larvae were used for larval bioassay testing. While 3-days sugar-fed adults derived from wild larvae after one generation under laboratory conditions were used for adult bioassay testing.

#### Insecticides

The insecticides used in this study for the larval and adult susceptibility tests were the diagnostic dosages as specified in the WHO standard methods.<sup>39,40</sup> The insecticides were obtained from WHO Collaborating Centre in

Malaysia (Vector Control Research Unit, School of Biological Sciences, Universiti Sains Malaysia, Penang). Larvae were tested against two Insect Growth Regulators (IGRs); Diflubenzuron (20 mg/l, 4 mg/l, 0.8 mg/l, 0.16 mg/l and 0.032 mg/l) and Methoprene (20 mg/l, 4 mg/l, 0.8 mg/l, 0.16 mg/l and 0.032 mg/l) and organophosphate; Temephos (0.005, 0.025, 0.125 and 0.625 mg/L). While adults were tested against organochlorine (DDT 4%), organophosphate (Fenitrothion 1%), four pyrethroids (Permethrin 0.75%, Lambda-cyhalothrin 0.05%, Deltamethrin 0.05%, Cyfluthrin 0.15%), and Carbamate (Bendiocarb 0.1%).

#### Larval Bioassay

Lots of 20 (third and fourth instar) larvae were placed in glass beakers filled with 249 ml of distilled water and 1 ml of each concentration of Diflubenzuron, Methoprene and Temephos. Five replicates of 20 larvae for each concentration and for control trials were used. For the Temephos, the numbers of dead larvae were recorded after 24 hours of the introduction of larvae to the beakers. While for the two IGRs, the duration of the observation period was that required for complete adult emergence in the control batches.

Percent inhibition of adult emergence (IE) was calculated and  $IE_{50}$  and  $IE_{95}$  values (effective concentration required for 50 and 95% emergence inhibition) were calculated for each of Diflubenzuron and Methoprene. Whereas, lethal concentration (LC) was calculated for Temephos ( $LC_{50}$  and  $LC_{95}$ ).

#### Adult bioassay

Sugar-fed, 3-5 days old-female mosquitoes were tested. A batch of 25 adults was introduced into holding tube before being exposed to insecticide-impregnated papers. Equal numbers of control tests were also carried out by exposing mosquitoes to insecticides-free papers. The experiment was replicated four times. After the respective period of exposure, all mosquitoes were transferred to new tubes, provided with 10% sugar solution and held for 24 hours recovery period. Mortality was recorded and resistance status was determined as per WHO criteria<sup>28</sup>; a population is considered susceptible if the mortality rate is 98-100%, the possibility of resistance (80-97%) and resistant (<80%).

#### Data analysis

Data were analyzed using computerized probit analysis **program** (Biostat 5) to determine the  $LC_{50}$  -  $LC_{95}$ , and  $IE_{50}$  -  $IE_{95}$  in the larval bioassay.

## Results and Discussion

### Larval susceptibility

The efficacy of Methoprene and Diflubenzuron against *A. aegypti* larvae is shown (Table-1). The result shows a high resistance of *A. aegypti* larvae to Temephos ( $LC_{50} = 61.8$  mg/l., and the best percentage mortality was obtained at the highest diagnostic dose of 0.625 mg/l was only 10%). Prolonged use of Temephos as a larvicide in Jazan region since 1986 could be one possible reason. Resistance to Temephos was detected in the region at 2003. Many authors around the world have reported resistance to Temephos in Havana<sup>20</sup>, Costa Rica<sup>5</sup>, Brazil<sup>6,17,19</sup>, and Thailand<sup>12</sup>. Similar results were also recorded in Bolivia and Argentina.<sup>4,27</sup> Contradictory, these findings are in contrast to the results reported in India<sup>31</sup>, where *A. aegypti* larvae were found susceptible to the Temephos. While in Malaysia, *A. aegypti* larvae showed to be tolerant to Temephos.<sup>9,16</sup>

The development of a new class of synthetic insect growth regulators (IGRs) Benzoylphenylureas (Chitin synthesis inhibitors) was a successful step forward towards non-hazardous and eco-safe Integrated Pest management (IPM).<sup>34</sup>

The results of the efficacy of two IGRs, Diflubenzuron and Methoprene, against *A. aegypti* larvae are shown (Table-2).

At the maximum dose of 20 mg/l, Methoprene and Diflubenzuron produced (100% and 87.5%) mortality against *A. aegypti* larvae, respectively. The  $IE_{50}$  of the Methoprene and Diflubenzuron were 0.49 and 0.86 mg/l, respectively. This result indicates that mosquito larvae of *A. aegypti* were more susceptible to Methoprene than Diflubenzuron by 1.8 folds.

The efficacy of Diflubenzuron against *A. aegypti* in this study is in contradiction with the

earlier<sup>23,28</sup> who noted that the Diflubenzuron was highly effective for inhibition of adult emergence, and (100%) adult emergence inhibition was observed in the presence of 0.1 ppm.

With low concentration (0.032 mg/l), Methoprene was found to be less effective in reducing the emergence of adult *A. aegypti* mosquito (Table 2), a result that matches the findings earlier.<sup>22</sup>

The  $IE_{50}$ -  $IE_{95}$  range (0.86 – 93.8 mg/l) post exposure (Table-2) against Diflubenzuron indicated less effect at larval stages as compared with Methoprene (0.49 - 10.9 mg/l). This finding of the effectiveness of the Methoprene against *A. aegypti* larvae in Jazan region is more relevant to the results obtained by other workers.<sup>7,8,10,23,30,33</sup>

In Malaysia, it was found that the larvae of *A. aegypti* exhibited moderate resistance toward Methoprene and low resistance toward Pyriproxyfen, with resistance ratios of 12.7 and 1.4, respectively, but susceptibility to Diflubenzuron.<sup>15</sup>

In this study, comparing the larvicidal activity of Methoprene with Diflubenzuron and Temephos (Tables- 1 & 2) it can be inferred that Methoprene is more effective against *A. aegypti* and can be used as an alternative larvicide in the areas where the low efficacy of Diflubenzuron and high resistance to organophosphate compounds have been detected.

### Adult susceptibility

The mortality of *Aedes aegypti* 24 hours after exposure to Deltamethrin, Permethrin, Cyfluthrin, Lambda-cyhalothrin and DDT is presented in Table-3.

The adult susceptibility from the different parts of Jazan revealed that the mosquitoes were susceptible to Cyfluthrin (100% mortality) and having a possibility of resistance to each of Fenitrothion, Deltamethrin, and Permethrin (93.6%,

**TABLE- 2 : Efficacy of Diflubenzuron and Methoprene against *A. aegypti* larvae**

Insecticides (IGRs)	Concentration (mg/l)	Larval Mortality (%)	* $IE_{50}$ ( $IE_{95}$ ) (mg/l)	95% Confidence limit	
				Lower $IE_{50}$ ( $IE_{95}$ )	Upper $IE_{50}$ ( $IE_{95}$ )
Diflubenzuron	0.032 - 20	17 - 87.5	0.86 (93.8)	0.48 (30.9)	1.52 (583.1)
Methoprene	0.032 - 20	18 - 100	0.49 (10.9)	0.0002 (1.49)	81.9 (7.57)

\*IE = Inhibition of adult Emergence

93.3%, and 86.8%, respectively). While, mosquitoes were resistant to Lambda-cyhalothrin (76.5%) and DDT (56.3%), as well as, they were highly resistant to Bendiocarb (17%).

It can be seen that *A. aegypti* showed the possibility of resistance to Deltamethrin and Permethrin (mortality 93.3%, 86.8%, respectively) in Jazan region. This result is in accordance with the findings of other authors in Thailand who reported the resistance of *A. aegypti* to Permethrin and Deltamethrin<sup>13</sup>. Whereas, the possibility of resistance of *A. aegypti* to the two insecticides is contradicted with the work of authors in India who found that adults *Aedes aegypti* and *A. albopictus* were susceptible to Deltamethrin and Permethrin<sup>29</sup>.

Deltamethrin and Permethrin could have resulted from cross-resistance with DDT. Moreover, the use of household aerosol insecticides where the main active ingredient is Permethrin might have contributed to the resistance of the species.

The development of high resistance against Bendiocarb (17%) and possibility of resistance against Fenitrothion (93.6%) could be due to the selection for resistance in the mosquitoes resulting from municipal and agricultural applications.<sup>18</sup> In addition, Fenitrothion has been used in the region as an interdomiliary residual spray between 1987

and 1994. However, the high resistance to Bendiocarb in our study is contrasting the findings of workers in Costa Rica, where *A. aegypti* was susceptible to it.<sup>5</sup>

In this study, *A. aegypti* mosquitoes showed resistance to Lambda-cyhalothrin and DDT, a result which was similarly obtained by other authors in India<sup>29</sup> with the exception only to the susceptibility of *A. aegypti* to Lambda-cyhalothrin. Conversely, when taking the LC<sub>50</sub> into consideration, workers in Jeddah found the larvae of *A. aegypti* to be susceptible to Lambda-cyhalothrin at a concentration of 0.01 ppm.<sup>2</sup> In Jazan region, the mosquito larvae have exhibited resistance to DDT as early as 1987 where the insecticide has long been used to control the vector of malaria.

Our findings show the appearance of resistance to multiple insecticides (e.g. Pyrethroids and Organochlorines) in *A. aegypti*. The same findings of multiple resistances between Pyrethroids and another group (Organophosphates) have been reported in South-East Asia<sup>13</sup>, South America and the Caribbean.<sup>21,25</sup>

The extensively successive use of Pyrethroids (Lambda-cyhalothrin, Deltamethrin, and Permethrin) in the region to control Malaria, Dengue and Rift valley mosquito vectors by the

**TABLE-3 : Mortality of adult *Aedes aegypti* 24 Hours after Exposure to Deltamethrin, Permethrin, Cyfluthrin, Lambda-cyhalothrin, Fenitrothion, Bendiocarb, and DDT.**

Insecticide	Concentration (%)	Exposure period (Hours)	Number of Mosquitoes Tested	Mortality (%)	Resistance status
Deltamethrin	0.05	1	120	93.3	Possibility of resistance
Permethrin	0.75	1	120	86.8	Possibility of resistance
Cyfluthrin	0.15	1	120	100	Susceptible
Lambda-cyhalothrin	0.05	1	120	76.5	Resistant
DDT	4	1	120	56.3	Resistant
Fenitrothion	1	1	100	93.6	Possibility of resistance
Bendiocarb	0.1	1	100	17	Resistant

Ministries of Health, Municipality and Agriculture, respectively, might have been contributed to accelerating the resistance to this group of insecticides.

This study revealed the effectiveness of Cyfluthrin against the adults of *A. aegypti* (100% mortality). This result may be attributed to the fact that Cyfluthrin is newly introduced to the region.

The resistance of *A. aegypti* against the tested insecticides decreased in the order of:

Bendiocarb > DDT > Lambda-cyhalothrin > Permethrin > Deltamethrin > Fenitrothion > Cyfluthrin.

### Conclusion

In this study, Cyfluthrin was found to be the only effective Pyrethroid used in Jazan region against the adult *A. aegypti*. Methoprene on the other hand proved to be an effective IGR against the larvae of *A. aegypti* in the region.

These results suggest that continuous resistance monitoring should be conducted regularly to identify the efficacy of compounds for dengue control. Well-managed rotation of the effective insecticides, community awareness and participation and public health campaigns to reduce larval breeding sites are recommended strategies to control dengue infections.

Moreover, the use of IGRs should be considered as an alternative when the larvae of *A. aegypti* develop resistance to conventional insecticides. Further studies should be carried out to provide more information about resistance status and to identify the resistance mechanism by PCR test in order to confirm and supplement the bioassay results.

### Conflict of interest

We declare that we have no conflict of interest.

### References

1. ADELA. AL-SHEIKH ( 2011) Larval habitat Ecology, Seasonal abundance and vectorial role in malaria transmission of *Anophelese arabiensis* in Jazan region of Saudi Arabia. *Journal of the Egyptian Society of Parasitology*,; **41**(3): 615-634.
2. AL-GHAMDI K, Z.I. AL-FIFI, M.S.SALEH, H.A. AL-QHTANI AND J.A. MAHYOUB (2008) Insecticide susceptibility of *A.aegypti*, the vector of dengue fever, in Jeddah governorate, Saudi Arabia. *Biosciences,Biotechnology Research Asia*,; **5**(2) : 501-506.
3. AL THBIANI AZIZ, HAMADY DIENG, AHMAD ABU HASSAN, TOMOMITSU SATHO, FUMIO MIAKE, MD RAWI CHE SALMAH AND SAZALY ABUBAKAR (2011) Insecticide susceptibility of the dengue vector *A.aegypti* ( Diptera: culicidae) in Makkah City,Saudi Arabia. *Asian Pacific Journal of Tropical Disease* 06/; **1**(2):94-99.
4. BIBER, P., RONDAN-DUEN, A.S. J., DUDUEN, A. A.F., GARDENAL, C. AND ALMIRON, W. (2006) Laboratory evaluation of susceptibility of natural subpopulations of *Aedes aegypti* larvae to temephos. *J Am Mosq Control Assoc*,; **22**: 408–411.
5. BISSET, J.A., MARÍN, R., RODRÍGUEZ, M.M., SEVERSON, D.W., RICARDO, Y., FRENCH, L., DÍAZ, M. AND PÉREZ, O. (2013) Insecticide resistance in two *Aedes aegypti* (Diptera: Culicidae) strains from Costa Rica. *J Med Entomol*,; **50**(2):352-61.
6. BRAGA, I.A., LIMA, J.B.P., SOARES, S.S. AND VALLE, D. ( 2004) *Aedes aegypti* resistance to temephos during 2001 in several municipalities in the states of Rio de Janeiro, Sergipe and Alagoas, Brazil. *Mem Inst Oswaldo Cruz*,; **99**: 199–203.
7. BRAGA, I.A., MELLO, C.B., MONTELLA, I.R., LIMA, J.B., MARTINS, ADE. J., MEDEIROS, P.F. AND VALLE, D. (2005) Effectiveness of methoprene, an insect growth regulator, against temephos-resistant *Aedes aegypti* populations from different Brazilian localities, under laboratory conditions. *J Med Entomol*. **42**(5):830-7.
8. CHEN, C.D., SELEENA, B., CHIANG, Y.F. AND LEE, H.L. (2008) Field evaluation of the bioefficacy of diflubenzuron (Dimilin®) against container-breeding *Aedes* sp. mosquitoes. *Tropical Biomedicine*,; **25**(1): 80–86 .
9. CHEN, C.-D., W. A. NAZNI, H.-L. LEE, AND SOFIAN-AZIRUN, M. (2005) Susceptibility of *Aedes aegypti* and *Aedes albopictus* to temephos in four sites in Kuala Lumpur City Center and Selangor

- State, Malaysia. *Trop. Biomed.*; **22**: 207–216.
10. GARG, R.C. AND W.A. DONAHUE (1989) Pharmacologic profile of methoprene, an insect growth regulator, in cattle, dogs, and cats. *J. Am. Vet. Med. Assoc.*; **194**: 410-412. *Hyg* **68**:329–333.
  11. GUBLER, D.J. (2002) Epidemic dengue/dengue hemorrhagic fever as a public health, social and economic problem in the 21st century. *Trends Microbiol.* **10**, 100–103. doi:10.1016/S0966-842X(01)02288-0.
  12. JIRAKANJANAKIT, N., SAENGTHARATIP, S., RONGNOPARUT, P., DUCHON, S., BELLEC, C., AND YOKSAN, S. (2007) Trend of Temephos resistance in *Aedes* (*Stegomyia*) mosquitoes in Thailand during 2003-2005. *Environ Entomol.* **36**, 506–511. Doi: 10.1603/0046-225X**36** [506: TOTRIA] 2.0.CO; 2.
  13. JIRAKANJANAKIT, N., RONGNOPARUT, P., SAENGTHARATIP, S., CHAREONVIRIYAPHAP, T., DUCHON, S., BELLEC, C. AND YOKSAN, S. (2007) Insecticide susceptible/resistance status in *Aedes* (*Stegomyia*) *aegypti* and *Aedes* (*Stegomyia*) *albopictus* (Diptera: Culicidae) in Thailand during 2003-2005. *J Econ Entomol.* **100**, 545–550. Doi: 10.1603/0022 0493**100**[545: IRSIAS] 2.0.CO; 2.
  14. KOW, C.Y., KOON, L.L. AND PANG, F.Y. (2001) Detection of dengue viruses in field caught male *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae) in Singapore by type-specific PCR. *J Med Entomol.*; **38**, 475–479. Doi: 10.1603/0022- 2585-38.4.475.
  15. LAU, K.W., CHEN, C.D., LEE, H.L., NORMA-RASHID, Y. AND SOFIAN-AZIRUN, M. (2015) Evaluation of insect growth regulators against field-collected *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae) from Malaysia. *Journal of medical entomology.* Mar 1; **52**(2):199-206.
  16. LEE, H.-L., AND W. LIME (1989) A re-evaluation of the susceptibility of field collected *Aedes* (*Stegomyia*) *aegypti* (Linnaeus) larvae to temephos in Malaysia. *Mosq. Borne Dis. Bull.* **4**: 91–95.
  17. LIMA, J.B.P., PEREIRA, D.A., CUNHAM, SILVA RCS, GALARDO, A.K.R., SOARES, S.S., BRAGA, I.A., RAMOS, R.P. AND VALLE, D. (2003) Resistance of *Aedes aegypti* to organophosphates in several municipalities in the states of Rio de Janeiro and Espirito Santo, Brazil. *Am J Trop Med Hyg.*; **68**: 329-333.
  18. LINES, J.D. (1988) Do agricultural insecticides select for insecticide resistance in mosquitoes? A look at the evidence. *Parasitol Today.*; **4**(7):S17-20.
  19. MACORIS, M.L.G., ANDRIGHETTI, M.T.M., TAKAKU, L., GLASSER, C.M., GARBELOTO, V.C. AND BRACCO, J.C. (2003) Resistance of *Aedes aegypti* from the state of S. Paulo, Brazil, to organophosphates insecticides. *Mem Inst Oswaldo Cruz.* **98**:703–708.
  20. MAGDALENA, RODRÍGUEZ, M., BISSET, J.A., FERNÁNDEZ, D. AND Omayda, P. (2004) Resistance to insecticides in larvae and adults of *Aedes aegypti*, Havana City: prevalence of A4 esterase associated with resistance to temephos. *Rev Cubana Med Trop.*; **56**(1):54-60.
  21. MARCOMBE, S., CARRON, A., DARRIET, F., ETIENNE, M., AGNEW, P., TOLOSA, M., MMY P- TCHA, LAGNEAU, C., YÉBAKIMA, A. AND CORBEL, V. (2009) Reduced Efficacy of Pyrethroid Space Sprays for Dengue Control in an Area of Martinique with Pyrethroid Resistance. *Am J Trop Med Hyg.* **80**, 745–751.
  22. NAYAR, J.K., ALI, A. AND ZAIM, M. (2002) Effectiveness and residual activity comparison of granular formulations of insect growth regulators Pripoxifen and s-Methoprene against Florida mosquitoes in laboratory and outdoor conditions. *J. Am. Mosq. Contr. Assoc.*, **18**: 196-201.
  23. NUSRAT JAHAN, JAWEERIA RAZAQ AND AYESHA JAN. (2011) Laboratory Evaluation of Chitin Synthesis Inhibitors (Diflubenzuron and Buprofezin) Against *Aedes aegypti* Larvae from Lahore, Pakistan. *Pakistan J. Zool.*, **43**(6), pp. 1079-1084.
  24. PEYREFITTE, C., ROUSSET, D., PASTORINO, B., POUILLOT, R., BESSAUD, M., TOCK, F., MANSARAY, H., MERLE, O., PASCUAL, A., PAUPY, C., VESSIERE, A., IMBERT, P., TCHENDJOU, P., DURAND, J.P., TOLOU, H. AND GRANDADAM, M (2007) Chikungunya virus, Cameroon. *Emerg Infect Dis.*; **13**,

- 138 A.A. ALSHEIKH, W.S. MOHAMMED, E.M. NOURELDIN, O.M. DAFFALLA, K.J. SHRWANI, Y.A. HOBANI, F.A. ALSHEIKH, M.H. ALZHRANI & A.A. BINSAEED  
768–771.
25. RAWLINS, S.C., MARTINEZ, R., WILTSHIRE, S. AND LEGALL, G. (1998) A comparison of surveillance systems for the dengue vector *Aedes aegypti* in Port of Spain, Trinidad. *J Am Mosq Control Assoc.*; **14**, 131–136.
  26. SATHANTRIPHOP, S., PAEPORN, P. AND SUPAPHATHOM, K. (2006) Detection of insecticides resistance status in *Culex quinquefasciatus* and *Aedes aegypti* to four major groups of insecticides. *Tropical Biomedicine.*; **23**(1):97-101.
  27. SECCACINI, E., LUCY´, A. A., LICASTRO, S., ZERBA, E. AND MASUH, H. (2007) Resistencia a temefos de *Aedes aegypti* (Diptera: Culicidae) en localidades del norte argentino. *5th Regional Mosquito Meeting (Argentina). Rev Biología Acuática.*; **23**:48.
  28. SECCACINI, E., ALEJANDRO LUCIA, LAURA HARBURGUER, EDUARDO ZERBA, SUSANALICASTRO AND HECTOR MASUH (2008) Effectiveness of Pyriproxyfen and Diflubenzuron Formulations as Larvicides Against *Aedes aegypti*. *Journal of the American Mosquito Control Association.*; **24**(3):398–403.
  29. SHARMA, S.N., SAXENA, V,K. AND LAL, S. (2004) Study on susceptibility status in aquatic and adult stages of *Aedes aegypti* and *Ae. albopictus* against insecticides at international airports of south India. *The Journal of Communicable Diseases.*; **36**(3):177-81.
  30. SILVA, JULIANA, JUNQUERIRA, MENDES AND JÚLIO. (2007) Susceptibility of *Aedes aegypti* to the insect growth regulators diflubenzuron and methoprene in Uberlandia, State of Minas Gerais. *Sociedade Brasileira de Medicina Tropical-SBMT. 1/12/2007.*
  31. SINGH, R.K., HAQ, S., KUMAR, GAURAV, MITTAL, P.K. AND DHIMAN, R.C. (2013) Insecticide susceptibility status of dengue vectors *Aedes aegypti* and *Aedes albopictus* in India: a review *Dengue Bulletin*; **37**:177.
  32. SOMBOON, P., PRAPANTHADARA, L. AND SUWONKERD, W. (2003) Insecticide susceptibility test of *Anopheles minimus* s.1., *Aedes aegypti*, *Aedes albopictus*, and *Culex quinquefasciatus* in northern Thailand. *Southeast Asian J Trop Med Public Health.*; **34**(1): 87-93.
  33. SULAIMAN SALLEHUDIN; SITI HAJAR, A. S. AND HIDAYATULFATHI OTHMAN.(2004) Residual efficacy of insect growth regulators pyriproxyfen, triflumuron and smethoprene against *Aedes aegypti* (L.) in plastic containers in the field. *Tropical Biomedicine*; **21** (1) : 97-100. 9 ref. 21 1 97-100.
  34. TUNAZ, H. AND N. UYGUN. (2004) Insect growth regulators for insect pest control. *Turk.J.Agri.for.*; **28**:377-387.
  35. WHO (1981) *Instructions for determining the susceptibility or resistance of mosquito larvae to insecticides*. Geneva, World Health Organization.
  36. WHO (1981) *Instructions for determining the susceptibility or resistance of adult mosquitoes to organochlorine, organophosphate and carbamate insecticides – diagnostic test*. Geneva, World Health Organization.
  37. WHO (1992) *Vector resistance to pesticides. 15<sup>th</sup> report of the WHO Expert Committee on Vector Biology and Control*. Geneva, Switzerland: World Health Organization.
  38. WHO (2005) *Guidelines for laboratory and field testing of mosquito larvicides*. (Document WHO/CDS/WHOPES/GCDPP/13, Geneva, Switzerland, World Health Organization.
  39. WHO (2006) *Guidelines for testing mosquito adulticides for indoor residual spraying and treatment of mosquito nets*. (Document WHO/CDS/NTD/ WHOPES/GCDPP/3, Geneva, Switzerland, World Health Organization.
  40. WHO (2009) *Dengue: guidelines for diagnosis, treatment, prevention and control- New edition*. WHO/HTM/NTD/DEN.1
  41. YAICHAROEN, R., KIATFUENGFOO, R., CHAERONVIRIYAPHAP, T., AND RONGNOPARUT, P. (2005) Characterization of deltamethrin, resistance in field populations of *Aedes aegypti* in Thailand. *J Vect Ecol.* **30** : 144–150.