

WATER QUALITY OF PARICHHA DAM RESERVOIR IN RELATION TO FISHERIES IN JHANSI (U.P.) INDIA**NEEL RATAN¹, SHARAT SRIVASTAVA², ALOK PATHAK³
AND *U.N. SINGH¹**Department of Botany¹, Chemistry² and Zoology³,
D.V. (P.G.) College,
ORAI-285001 (U.P.) INDIA
*Corresponding Author:
Email : drunsingh@rediffmail.com**Received** : 04.03.17; **Revised** : 25.03.17; **Accepted** : 02.05.17**ABSTRACT**

Physico-chemical characteristics of Parichha dam reservoir of Jhansi in Uttar Pradesh have been studied. The water temperature varied between 23.0 to 37.0°C. The transparency, pH, chlorides and total hardness were in the range of 100 to 145 cm, 7.8 to 9.2, 42.0 to 59.17 mg/l and 100-197 mg/l respectively. The total alkalinity, dissolved oxygen and total dissolved solids ranged between 193 to 389, 5.2 to 8.6 and 90 to 310 mg/l respectively. The study revealed that the reservoir water is suitable for fisheries.

Figure : 00

References : 19

Tables : 02

KEY WORDS : Fisheries, Jhansi, Reservoir, Uttar Pradesh, Water quality.**Introduction**

The reservoir came into existence with impounding of Parichha dam in Jhansi (U.P.) India. The morphometric features are depicted (Table-1). The work on water quality of Parichha dam reservoir can provide valuable information to the planners, scientists and fishing authorities to check out strategies for the development of fisheries. There is meagre record of water quality of Parichha dam reservoir.

The study of physico-chemical characteristics of Parichha reservoir includes the water temperature, transparency, total dissolved solids, pH, chlorides, total hardness, dissolved oxygen, carbondioxide and total alkalinity.

Materials and Methods

The Parichha dam reservoir is situated at latitude 78° 36' E and longitude 25° 31' N at an elevation of approximately 258m above mean sea level in Bundelkhand region, district Jhansi, Uttar Pradesh.

The methods used for analysis of various physico-chemical parameters except pH are as given in methodology for water analysis¹⁸. The bottles used for collection of water samples were made up of Pyrex glass and were cleaned by boiling them in dilute HNO₃ followed by distilled water⁵. The characteristics like temperature, pH, conductivity, dissolved oxygen etc. were measured as early as possible after collection.

ACKNOWLEDGEMENTS : Authors are grateful to the Principal, D.V. Postgraduate College, Orai-285001 (U.P.) India for rendering laboratory facilities and encouragement to complete the present study. The authors also express their sincere thanks to the authorities of Parichha dam reservoir for providing morphometric information and kind permission to work with basic necessary assistance.

TABLE- 1: Morphometric features of Parichha dam reservoir in Jhansi district

1	River	Betwa
2	Location	Jhansi
3	Year of Construction	1881-86
4	Rainfall (mm)	989
5	Length of dam (m)	1174.69
6	Height of dam (m)	16.77
7	Water spread area (ha)	802
8	Irrigation potential (ha)	98500 (Approx.)

pH of the samples were determined by Equip-Tronics digital pH meter model EQ-610 using glass electrode. Specific conductance was measured by Equip-Tronics digital conductivity meter model EQ-660 A which is first calibrated by using standard KCl solution. Total dissolved solids was determined by standard method.¹ Total hardness and chlorids in the water samples were determined by literature with standard EDTA¹⁰. The dissolved oxygen in water samples was estimated by standard method.²⁰

Results and Discussion

The water quality data of reservoir is depicted (Table-2). Among all the ecological factors, the water temperature can be considered as an important factor on which the spawning in fishes depends. In the Indian major carps, the optimum temperature range for successful induction of spawning have been suggested from 24 to 31°C and beyond this range fish do not spawn⁶. As per observation in common carp, (*Cyprinus carpio*) gametogenesis may be completed in October but spawning does not occur until the water temperature rises above 17°C, the minimum spawning temperature³. The rate of chemical and biological reactions is being doubled with every 10°C increase in temperature. Warm water fish grow

the best at a temperature range⁴ (25 to 32°C). During the present study the minimum water temperature was recorded in November 2015 (23°C) and the maximum in April 2016 (37°C). Reservoirs having water temperature more than 22°C are found to be highly productive¹¹. The average water temperature during the present investigation was 27°C. Hence the reservoir water is highly productive.

Transparency is a physical variable significant to production. During the present study secchi disc transparency ranged from 100 to 145 cm. The minimum transparency value was recorded in July 2015 while the maximum in April 2016. Present findings collaborate with findings of others^{15,16} who reported the lowest transparency values in rainy season. The low values recorded during rainy season may be due to the heavy rains and winds of high velocity. Similarly high values during winter and summer period can be attributed to low moderate wind velocity. High secchi disc transparency values indicate low primary productivity by phytoplankton and consequent low fish production⁸.

The pH is affected not only by levels of carbon dioxide but also by other organic and inorganic components of water. Further, any alteration in water pH is accompanied by changes in other physico-chemical parameters. In the present study pH ranged from 7.8 to 9.2. The minimum value was recorded in May 2015 while the maximum in January 2016. The pH range between 6.0 to 8.5 indicates medium productive nature of a reservoir, more than 8.5 highly productive and less than 6 low productive reservoirs¹¹. In case of Parichha dam reservoir, the average pH value during the study period was observed to be 8.5 which indicated that the reservoir is a medium productive. It has been observed that change of pH may be one of the main factors which may induce the fish to spawn¹². On the other hand it is suggested that the interaction of many ecological factors like high oxygen content, relatively low free CO₂ and slightly alkaline pH were essential for the breeding of *Schizothorax*, *Labeo*, *Puntius ad Barilius spp.* of Garhwal Himalaya².

During present investigation the total dissolved solids were high in summer followed by winter and monsoon. The investigation of CIFRI⁷ also reported the range of total dissolved solids in Bhatghar reservoir of Maharashtra, being maximum in March and January, minimum in October and June at lentic and intermediate sectors of the

WATER QUALITY OF PARICHHA DAM RESERVOIR IN RELATION TO FISHERIES IN JHANSI (U.P.) INDIA 97

reservoir. In lotic sector it was maximum in December and minimum in July.

The dissolved oxygen is of great importance to all living organisms. It dissolves in water due to direct diffusion from air and/or photosynthetic activity of autotrophs. The diffusion of oxygen is dependent on temperature of water, its salinity, total dissolved solids, and movements of water **etc.** During the study period, the dissolved oxygen ranged from 5.2 to 8.6 mg/l. The high dissolved oxygen was found during monsoon, while lower values were recorded in summer.

Free carbon dioxide is an extremely necessary constituent in an aquatic environment¹⁹. Free carbon dioxide in water occurs due to respiration of aquatic biota, decomposition of organic matters and also due to infiltration through the soil. Carbon dioxide is an important component of the buffer system, which influences carbonate and bicarbonate concentrations in water. The fishes can tolerate a higher concentration of free carbon dioxide up to 60 mg/l, with a tendency to avoid as low⁹ as 5 mg/l⁹. In the present investigation, free carbon dioxide generally varied between nil and 2.5 mg/ L. It was not detected in January 2016 and April 2016.

Alkalinity of any water is mainly due to carbonates, bicarbonates and hydroxide. It is an index of nutrient status in a water body. The total alkalinity in Parichha dam reservoir ranged from 193 to 289 mg/l. The availability of carbon dioxide for primary production is related to alkalinity. Water with total alkalinity levels of 20 to 150 mg/L generally contains adequate quantity of carbon dioxide⁴. Water bodies having total alkalinity above 50 mg/l can be considered productive¹³. The total alkalinity observed¹⁴ was in accordance with the present finding.

The term hardness is frequently used as the assessment of the water quality. This is governed by the concentration of calcium and magnesium salts, which combined largely with bicarbonate, carbonate, sulphates, chlorides and

TABLE- 2: Water quality of Parichha dam reservoir in Jhansi

1.	Water temperature °C	23.0-37.0
2.	pH	7.8-9.2
3.	Transparency (cm)	100.0-145.0
4.	Total Dissolved Solids (mg/l)	90.0-310.0
5.	Chlorides (mg/l)	42.0-59.17
6.	Total hardness (mg/l)	100.0-197.0
7.	Dissolved oxygen (mg/l)	5.2-8.6
8.	Total alkalinity (mg/l)	193.0-289.0
9.	Free carbon dioxide (mg/l)	Nil-2.5

other anions of mineral acids. Desirable level of total hardness for fish culture generally fall within the range of 20.0 to 300.0 mg/L. It is suggested¹⁷ that a total hardness of 50.0 mg/L calcium carbonate equivalent is the dividing line between soft and hard waters and according to above suggestion pond waters having a total hardness of 15.0mg/L or above are good for fish growth¹⁷. Based on the average hardness level, Parichha dam reservoir can be considered suitable for fisheries. During the present investigation the maximum total hardness (197.0 mg/l) was found in May 2016, its minima (100.0 mg/l) was recorded in October 2015. The higher values of total hardness were recorded during summer and lower values were recorded in the winter season.

References

1. AWWA (1976) *Water Quality and Treatment*. Mc Graw Hill, New York.
2. BADOLA, S.P. AND SINGH, H.R. (1984) Spawning of some important coldwater fish of the Garhwal Himalaya. *Bombay Nat. Hist. Soc.* **81** (1) : 54-58.
3. BILLARD, R., BRETON, B., FOSTIER, A., JALABERT, B. AND WEIL, C. (1978) Endocrine control of the teleost reproductive cycle and its relation to external factors: samoid and cyprinid models. In:

- 98 **NEEL RATAN, SHARAT SRIVASTAVA, ALOK PATHAK AND *U.N. SINGH**
- Comparative Endocrinology*, (P.J. Gaillard and H.H. Boer eds.), Elsevier North-Holland Biomedical Press, Amsterdam, pp 37-48.
4. BOYD, C.E. (1984) *Water quality management for pond fish culture*. Elsevier Sci. Publ. Co., Amsterdam, 318pp.
 5. BROWN, E., SOUGSTAD, M.W. AND FISHMAN, M.J. (1974) *Method for Collection and Analysis of Water Samples for Dissolved Minerals and Gases*. U.S. Dept. of Interior. Book No. 5.
 6. CHAUDHARI, H. (1950) Experiments on induced spawning of Indian carps with pituitary injections. *Ind. J. Fish.* **7** (2) : 20-49.
 7. CIFRI (1997) Ecology and Fisheries of Bhatghar reservoir. *Golden Jubilee spl. Bulletin No. 73*.
 8. DAS, R.K. (1996) Monitoring of water quality, its importance in disease control. Paper presented In: *Nat. Workshop on Fish and Drawn Disease, Epizootics and Quarantine Adoption in India*. October 9, 1996. CICFKI. pp 51-55.
 9. HART, J.S. (1944) The circulation and respiratory tolerance of some Florida freshwater fishes. *Proc. Acad. Sci.* **7**: 221.-246.
 10. JEFFERY, G.H., SASSET, J., MENDHAM, J. AND DENNY, R.C. (1978) *Vogel's Text Book of Quantitative Chemical Analysis*. 5th Ed. Wesley Longman Limited, England.
 11. JHINGRAN, A.G. AND SUGUNAN, V.V. (1990) General guidelines and planning criteria for small reservoir fisheries management. p.18. In: Jhingran Arun G. and V.K. Unnithan (eds). *Reservoir Fisheries in India. Proc. of the Nat. Workshop on Reservoir Fisheries*, 3-4 January 1990. Special Publication 3, Asian Fisheries Society, Indian Branch, Mangalore, India.
 12. MOOKERJEE, H.K. (1945) Factors influencing the spawning of principal carps in India. Symposium on The factors influencing the spawning of Indian carps. *Proc. Nat. Inst. Sci. India.* **71**: 312-315.
 13. MOYLE, J.B. (1949) Some indices of lake productivity. *Trans. Amer. Fish. Soc.* **76** : 322-334.
 14. SAKHARE, V.B. AND JETITHOR, S.G. (2009) Physico-chemical limnology of Pathri reservoir (Maharashtra) in relation to fisheries. In: *Aquatic Biology and Aquaculture* (Ed.Y.B. Sakhare), Manglam Publications, Delhi, pp 147-163.
 15. SAKHARE, V.B. AND JOSHI, P.K. (2002) Ecology of Palas Nilegaon reservoir in Osmanabad district, Maharashtra. *J. Aqua. Biol.* **18** (2) :17-22.
 16. SINGH, C.S., SHARMA, A.P. AND DEORARI, B.P. (1993) Plankton population in relation to fisheries in Nanak Sagar reservoir, Nainital. In: Prof. K.S. Rao (ed.). *Recent Advances in Freshwater Biology*. **1** : 66-79.
 17. SWINGLE, H.S. (1967) Standardization of chemical analysis of water and ponds muds. *FAO Fish. Rep.* **44** : 397-342.
 18. TRIVEDI, R.K. AND GOEL, P.K. (1986) Chemical and biological methods for water pollution studies. *Environmental Publication*, Karad, India.
 19. WELCH, T.S. (1952) *Limnology*, 2nd Ed. Mc Graw Hill Book Co. N.Y. 1952.