

## MOLLUSCAN DIVERSITY WITH PARTICULAR EMPHASIS TO SNAILS IN DISTRICT GAUTAM BUDH NAGAR (U.P.) INDIA

LOMESH KUMAR TYAGI

Department of Biotechnology,  
Lloyd Institute of Management & Technology,  
GREATER NOIDA (U.P.) 201306 INDIA  
Email: ltyagi\_98@rediffmail.com

*Received* : 18.08.2018; *Accepted* : 10.10.2018

### ABSTRACT

Present investigation reveals molluscan diversity of the 4 selected sites viz., Tilpata Karanbas, Manhcha, Latifpur and Atta Gujran in the vicinity of district Gautam Budh Nagar (U.P.). As many as 22 species of molluscs under 9 families were collected. There were 19 species of snails viz., *Bellamya bengalensis*, *B. dissimilis*, *B. crassa*, *Lymnaea (Pseudosuccinea) acuminata* f. *hians*, *L. luteola* f. *ovalis*, *L. acuminata* f. *patula*, *L. (Radix) auricularia*, *L. (Lymnaea) stagnalis*, *L. acuminata* f. *gracillior*, *Pila globosa*, *Indoplanorbis exustus*, *Gyraulus convexiculus*, *G. pankongensis*, *Planorbis planorbis* var. *tangitarenis*, *Hydrobioides nassa*, *Thiara (Melanoides) tuberculata*, *Thiara (Thiara) scabra*, *T. (M.) trigrina* and *Brotia (Antimelania) costula* under 6 families. The other molluscs were *Parreysia (Parreysia) favidens*, *Corbicula striatella* and *Lamellidens marginalis*. There has been a marked variation in the composition of molluscan species collected from different habitats.

Figures : 02

References : 23

Table : 01

KEY WORDS: Gautam Budh Nagar, Molluscan diversity, Snails, U.P.

### Introduction

Over 90,000 species of gastropods have been recorded worldwide, both in the water and on land. Although molluscs are common components of the benthic communities, their role in the dynamics of the aquatic ecosystem and their contribution to biomass production is not well known. Freshwater gastropods are either herbivores or detritivores, or they may passively consume small invertebrates associated with periphyton<sup>2</sup>.

Many species spend their entire lives in a few square meters of habitat, making them extremely vulnerable to localized environmental habitat degradation. Although most species prefer clean, stable and river bottoms, some prefer the soft substrates, more common to ponds and lakes. Besides this, a few wide-ranging snail species can easily survive in polluted habitats.

Our knowledge on Indian freshwater molluscs is based on the contributions made by several earlier workers<sup>22</sup> in the monograph "Freshwater molluscs of India" mentioned in brief about the general aspects like habitat distribution, Zoogeographical significance, importance of malacological studies, their role in medical veterinary, public health, aqua culture, etc. Some considerable work

on faunal diversity of snails of Uttaranchal (now Uttarakhand) region has been worked out in the past<sup>3,14,15,20,21</sup>. It is evident from the available literature that almost no work has been carried out on faunal diversity of snails from Gautam Budh Nagar, henceforth, the present effort is on molluscan diversity of the area so as to add additional information in the existing knowledge.

### Methodology

#### (A) Collection of the snails

The snails were collected twice a month from March 2012 to February 2014 using the articles such as hand or water scoop net, polythene bags, specimen's bottles, gloves and forceps. A standardized data-sheet was prepared pertaining to the snails. In each collection two persons were involved and one-hour time was spent in the collection of snails. The samples so collected were brought to the laboratory in the polythene bags and separated species wise. The snails were fixed in the Bouin's fluid for further studies.

#### (B) Identification of the snail's species

The snails were narcotized by providing a few drops of 70 % alcohol. Sometimes the chemicals like MnSO<sub>4</sub>,

**ACKNOWLEDGEMENT :** I am thankful to Dr. R.K. Jauhari, Head, Dept. of Zoology, D.A.V.(P.G.) College, Dehradun for useful suggestions in the preparation of manuscript.

Chloral hydrate, etc. were also used as narcotizing agents. The body of these animals was removed from the shell with the help of forceps and then the empty shells were boiled in water so as to remove any soft part if stuck inside the shell. Thereafter, the shell was dried in the air under sunlight and kept in small specimen's tube depending upon the size of the snail's sample. Relevant information such as the size, date and time of collection was also provided with each sample. The genera of the snails were identified in the laboratory with the help of the Keys for identification of snails as given in Fauna of British India<sup>4</sup>, freshwater molluscs of India<sup>9</sup> and Handbook on Indian Freshwater Molluscs<sup>19</sup>. In case any confusion existed in the identification of the collected snails, the samples were sent to the Zoological Survey of India, Kolkata for confirmation of species. Further studies were continued only after getting the confirmation of the species from the concerned agency.

## Results

Survey work towards the molluscan diversity of the 4 selected sites viz., Tilpata Karanbas (L-1), Manhcha (L-2), Latifpur (L-3) and Atta Gujran (L-4) in the vicinity of district Gautam Budh Nagar (U.P.) between March 2012 – February 2014 revealed the occurrence of 22 species of molluscs under 9 families (Table-1 and Fig. 1). Of these, there were the following 19 species of snails viz., *Bellamyia bengalensis*; *B. dissimilis*; *B. crassa*; *Lymnaea (Pseudosuccinea) acuminata* f. *hians*; *L. luteola* f. *ovalis*; *L. acuminata* f. *patula*; *L. (Radix) auricularia*; *L. (Lymnaea) stagnalis*; *L. acuminata* f. *gracilior*; *Pila globosa*;

*Indoplanorbis exustus*; *Gyraulus convexciculus*; *G. pankongensis*; *Planorbis planorbis* var. *tangitarensis*; *Hydrobioides nassa*; *Thiara (Melanoides) tuberculata*; *Thiara (Thiara) scabra*, *T. (M.) trigrina* and *Brotia (Antimelania) costula* under 6 families. The other molluscs were *Parreysia (Parreysia) favidens*, *Corbicula striatella* and *Lamellidens marginalis*. Further, the commonly occurring snails were *Bellamyia bengalensis*, *L. luteola* f. *ovalis*, *L. acuminata* f. *patula*, *Pila globosa*, *Indoplanorbis exustus*, *Gyraulus convexciculus*, *Planorbis planorbis* var. *tangitarensis* and *Thiara (Melanoides) tuberculata*. The following 3 species viz., *L. (Lymnaea) stagnalis*, *B. crassa* and *Brotia (Antimelania) costula* have been recorded from Manhcha (L-2) locality. However, the species like *G. pankongensis*, has been recorded from L-3 locality. Among other molluscs, most commonly occurring species were *Lamellidens marginalis* followed by *Parreysia (Parreysia) favidens* and *Corbicula striatella* in succession.

There were 21 species of molluscs collected from Manhcha (L-2) locality while from the other 3 localities, the number was 18, 17 and 15 species, respectively from Latifpur (L-3), Atta Gujran (L-4) and Tilpata Karanbas (L-1). There has been a marked variation in the composition of molluscan species collected from different habitats. The details about the faunal diversity of molluscs along with the habitats in each locality have been presented separately.

### a) Tilpata Karanbas (L-1) locality

At this locality as many as 15 species of molluscs




Fig.1: Composition of Molluscan families in selected localities of district Gautam Budh Nagar during the study period (2012-2014)

TABLE-1: Molluscan diversity in selected localities of district Gautam Budh Nagar(U.P.) March 2012 to February 2014.

Sl. No.	Family	Molluscan species	Locality			
			Tilpata Kambas (L-1)	Manhcha (L-2)	Latifpur (L-3)	Atta Gujran (L-4)
1	Viviparidae	<i>Bellamyia bengalensis</i>	+	+	+	+
2		<i>B. dissimilis</i>	-	+	+	-
3		<i>B. crassa</i>	-	+	-	-
4	Lymnaeidae	<i>Lymnaea (Pseudosuccinea) acuminata f. hians</i>	+	+	+	+
5		<i>L. luteola f. ovalis</i>	+	+	+	+
6		<i>L. acuminata f. patula</i>	+	+	+	+
7		<i>L. (Radix) auricularia</i>	+	+	+	+
8		<i>L. (Lymnaea) stagnalis</i>	-	+	-	-
9		<i>L. acuminata f. gracilior</i>	+	+	+	+
10	Pilidae	<i>Pila globosa</i>	+	+	+	+
11	Planorbidae	<i>Indoplanorbis exustus</i>	+	+	+	+
12		<i>Gyraulus convexiculus</i>	+	+	+	+
13		<i>G. pankongensis</i>	-	-	+	+
14		<i>Planorbis planorbis var. tangitarenis</i>	+	+	+	+
15	Bithyniidae	<i>Hydrobioides nassa</i>	-	+	+	+
16	Thiaridae	<i>Thiara (Melanoides) tuberculata</i>	+	+	+	+
17		<i>Thiara (Thiara) scabra</i>	+	+	-	-
18		<i>T. (M.) trigrina</i>	+	+	+	+
19		<i>Brotia (Antimelania) costula</i>	-	+	-	-
20	Amblemidae	<i>Parreysia (Parreysia) favidens</i>	+	+	+	+
21	Corbicullidae	<i>Corbicula striatella</i>	-	+	+	+
22	Unionidae	<i>Lamellidens marginalis</i>	+	+	+	+

under 7 families have been recorded from 5 different habitats like streams, river beds, ponds, rice fields and tanks during 2012-2013. The maximum number of specimens of molluscs was recorded from river streams while the species diversity was comparatively more in the river beds. Less number of specimens was recorded from tanks. Rice fields also contributed to the molluscan diversity to a great extent. Commonly occurring snail's species were *Bellamyia bengalensis*, *Lymnaea (Pseudosuccinea) acuminata* f. *hians*, *L. luteola* f. *ovalis*, *L. acuminata* f. *patula*, *L. (Radix) auricularia*, *Indoplanorbis exustus* and *Planorbis planorbis* var. *tangitarenis*. Further, *L. acuminata* f. *gracilior* has been recorded from river beds and rice fields. Similarly, *Lamellidens marginalis* has been found in river beds. Almost the same kind of pattern about the number of species, their occurrence in various habitats has been recorded in the year 2013-2014 (Table-1)

#### b) Manhcha (L-2) locality

During 2013-2014, at Manhcha locality as many as 21 species of molluscs under 9 families have been recorded from 5 different habitats like streams, river beds, ponds, rice fields and tanks. The maximum number of specimens of molluscs was recorded from rice fields followed by river beds and ponds in succession. Further, the species diversity was comparatively more in the rice fields. Less number of specimens was recorded from tanks. River beds contributed to the molluscan diversity to a great extent. Commonly occurring snail's species were *Bellamyia bengalensis*, *Lymnaea (Pseudosuccinea) acuminata* f. *hians*, *L. luteola* f. *ovalis*, *L. acuminata* f. *patula*, *L. (Radix) auricularia*, *Indoplanorbis exustus*, *Thiara (Melanoides) tuberculata* and *Planorbis planorbis* var. *tangitarenis*. Further, *Hydrobioides nassa* has been recorded from rice fields. *Corbicula striatella* and *Lamellidens marginalis* have been found in river beds, streams and rice fields. Almost the same kind of pattern about the number of species, their occurrence in various habitats has been recorded in the year 2013-2014. However, the population of molluscs was comparatively more in 2013-14 than 2012-13 (Table-1 & Fig. 1)

#### c) Latifpur (L-3) locality

As many as 18 species of molluscs under 9 families have been recorded from the 5 different habitats i.e., marshy land, ponds, streams, rice fields and tanks during 2012-2014. The maximum number of specimens of molluscs was recorded from rice fields. Further, species diversity was comparatively more in the rice fields followed by streams / marshy land, ponds and tanks in succession during 2012-2013. Like L-1 and L-2 localities, less number of specimens was recorded from tanks. Commonly occurring snail's species from L-3 locality were *Bellamyia*

*bengalensis*, *Lymnaea (Pseudosuccinea) acuminata* f. *hians*, *L. luteola* f. *ovalis*, *L. acuminata* f. *patula*, *L. (Radix) auricularia*, *Indoplanorbis exustus*, *Planorbis planorbis* var. *tangitarenis*, *Thiara (Melanoides) tuberculata* and *Corbicula striatella*. Further, *T. (M.) trigrina* has been recorded from marshy land and rice fields. Similarly, *B. dissimilis* and *Hydrobioides nassa* have been found in rice fields and streams while *Gyraulus convexiculus* has been recorded from rice fields. Although almost the same kind of pattern about the number of species, their occurrence in various habitats has been recorded in the year 2013-2014. In fact, the population of molluscs was more in 2013-14 (Table-1 & Fig. 1)


#### d) Atta Gujran (L-4) locality

During 2013-2014, at Atta Gujran locality as many as 17 species of molluscs under 9 families have been recorded from 4 different habitats like, marshy land, ponds, rice fields and tanks. The maximum number of specimens of molluscs was recorded from rice fields followed by streams / ponds and marshy land in succession. Further, the species diversity was comparatively less in the tanks. Commonly occurring snail's species were *Bellamyia bengalensis*, *Lymnaea (Pseudosuccinea) acuminata* f. *hians*, *L. luteola* f. *ovalis*, *L. acuminata* f. *patula*, *L. (Radix) auricularia*, *Indoplanorbis exustus*, *Thiara (Melanoides) tuberculata*, *G. pankongensis* and *Planorbis planorbis* var. *tangitarenis*. Further, *L. acuminata* f. *gracilior* has been recorded from rice fields and streams. *Corbicula striatella* and *Lamellidens marginalis* have been found to occur in almost all the habitats. *Hydrobioides nassa* has been recorded from ponds, streams and rice fields. Almost the same kind of pattern about the number of species, their occurrence in various habitats has been recorded in the year 2013-2014 (Table-1 & Fig. 1).

### Discussion

The present findings reveal occurrence of 22 species of molluscs under 9 families. Of these, there were 19 species of snails viz., *Bellamyia bengalensis*; *B. dissimilis*; *B. crassa*; *Lymnaea (Pseudosuccinea) acuminata* f. *hians*; *L. luteola* f. *ovalis*; *L. acuminata* f. *patula*; *L. (Radix) auricularia*; *L. (Lymnaea) stagnalis*; *L. acuminata* f. *gracilior*; *Pila globosa*; *Indoplanorbis exustus*; *Gyraulus convexiculus*; *G. pankongensis*; *Planorbis planorbis* var. *tangitarenis*; *Hydrobioides nassa*; *Thiara (Melanoides) tuberculata*; *Thiara (Thiara) scabra*; *T. (M.) trigrina* and *Brotia (Antimelania) costula* under 6 families. There are a number of workers who conducted studies on molluscan faunal diversity from different parts of India and almost similar findings were reported<sup>1,5-8,12</sup>.

The occurrence of *Bellamyia bengalensis*



**Fig. 2: Species composition of collected molluscs at selected localities of district Gautam Budh Nagar during the study period (2012-2014) .**

(Lamarck), *Melanoides tuberculatus*, *M. scabra*, *Lymnaea acuminata*, *L. luteola*, *Indoplanorbis exustus* and *Pila* sp. during the present is directly supported by the findings<sup>2,12</sup> but they also reported one or more species of molluscs.

From Gwalior (M.P.), a brief account on 9 species of freshwater snails including *Indoplanorbis exustus*, *Lymnaea (Pseudosuccinea) luteola*, *L. (P.) acuminata f. rufescens* and *Vivipara bengalensis* together with notes on their habitats and a list of trematode larvae was furnished<sup>14</sup>. There is a difference with in respect of both the species of snails and the trematode larvae.

The present findings also differ with earlier<sup>12</sup> which reported 14 species of Gastropods and Bivalves from the irrigated area of Surat district (Gujarat). A study on the molluscan diversity of Asan river system in Doon Valley with special reference to vectors of trematode parasites collected 18 species of molluscs comprising 13 species of gastropods grouped under 5 different families<sup>14</sup>. The results of the present study are in accordance with<sup>14</sup> in respect of occurrence of certain common forms like *Bellamya bengalensis*, *L. luteola f. ovalis*, *L. acuminata f. patula*, *Pila globosa*, *Indoplanorbis exustus*, *Gyraulus convexisculus*, *Planorbis planorbis var. tangitarenis* and *Thiara (Melanoides) tuberculata*.

In a study on the molluscan fauna of Narmada Sagar were collected eleven species of molluscs, out of which 8 species comprises of *Rachis bengalensis*, *R. punctatus*, *Bellamya bengalensis*, *Melanoides tuberculatus*, *M. scabra*, *Lymnaea acuminata*, *L. luteola* and *Indoplanorbis exustus*. The species of mollusks as gathered in the present study were quite different as reported earlier<sup>2</sup>.

A study on the molluscan diversity of Saipung Wildlife Sanctuary, Meghalaya revealed 13 species of molluscs, out of which 12 species were identified as gastropods and one species of Bivalve<sup>10</sup>. The freshwater gastropod species comprises *Bellamya bengalensis f. annadalei*, *Pila theobaldi*, *Thiara (Iareba) lineata*, *Brotia (Antimelania) costula*, *Paludomus (Paludomus) conica*, *P. (P.) regulata*, *P. (P.) stephanus* and *Indoplanorbis exustus*. There is a difference between the findings of present study and the earlier one<sup>10</sup>.

Keeping in view the above statement, it is clear that diversity exists in the species of the snails. This may be due to different reasons in the fluctuating pattern of the diversity. Further, either a single factor alone or a combination of factors is responsible for such variation.

## References

1. Agrawal HP, Gupta PD. Mollusc fauna of Narmada Sagar and impact assessment of the proposed dam. *Cheetal*, 1994; **33** (2): 20-29.
2. Cuker BE. Competition and coexistence among the grazing snail *Lymnaea*, Chironomidae, and microcrustacea in an arctic epilithic lacustrine community. *Ecol.* 1983; **64** : 10-15.
3. Goel HC, Srivastava CP. Freshwater snails of Gwalior (M.P.). *J. Bomb. Nat. Hist. Soc.* 1978; **77** : 215-222
4. Gupta SD. Macrobenthic fauna of Loni Reservoir. *J. Inland Fish. Soc. India.* 1976; **8** : 49-59.
5. Hatter SJS, Sen N, Mathew R, Sharma S. Faunal diversity of Saipung Wild Life sanctuary, Meghalaya. 2004; 1-6.

6. Mitra SC, Dey A, Ramakrishna. Freshwater molluscan fauna of Asan wetland, Wetland ecosystem series. *Zool. Surv. India*. 2003; **5** : 19-22.
7. Pandey AP, Mehta NR. Freshwater snails of irrigated area of Surat District, Gujarat, India. *J. Zool. Soc. India*. 1982; **34** (1&2) : 118-120
8. Patil SG, Ramakrishna. Fresh water mollusks Fauna of Nathsagar wet land. Wet land Ecosystem series 7. *Zool. Surv. India*. 2005; 101-114
9. Patil SG. Fresh water molluscs: Fauna of Melghat Tiger Reserve, *Conservation area series, 24*, *Zool. Surv. India*. 2005; 437-449
10. Pokhriyal BP, Mahesh RK, Jauhari RK. Prevalence of trematode cercarial infection in the snail *Lymnaea (Pseudosuccinea) acuminata* Lamarck, 1822 in different localities of Dehra Dun Valley. *Bioved.* 1997; **8** (1&2): 15-19.
11. Pokhriyal BP, Mahesh RK, Jauhari RK. Studies on the molluscan diversity in Asan river system in Doon valley with special reference to vectors of trematode parasites. *Indian J. Forestry*. 1998a; **21** (2) : 167-170.
12. Pokhriyal BP, Mahesh RK, Jauhari RK. Occurrence of recognized helminthic vector snails in different habitats in Dehradun Valley. *Him. J. Env. Zool.* 1998b; **12** : 167-172.
13. Preston HB. Fauna of British India including Ceylon and Burma, Mollusca (Freshwater Gastropoda and Pelecypoda), London. xix+ 1915; 244 pp.
14. Ramakrishna, Dey A. Handbook on Indian freshwater Molluscs, Zoological Survey of India. 2007; 399p.
15. Ray HC, Mukherjee A. Mollusca fauna of Rajasthan, India. *Rec. Zool. Surv. India*. 1963; 403-433
16. Srivastava N, Jauhari R. Population fluctuations of the freshwater snail *Thiara (Melanoides) tuberculata* in Doon Valley in relation to certain factors. *J. Zool. Soc. India*. 1990a; **41** (1&2) : 63-67.
17. Srivastava N, Jauhari RK. Influence of certain abiotic factors on the occurrence of larval trematodes in *Lymnaea acuminata* (Lamarck) in Doon Valley. *Riv. di Parasit.* 1990b; **6** (1): 335-341.
18. Subba Rao NV. Freshwater Molluscs of India. In: Roa K.S. (Ed.). *Recent Advances in Freshwater Biology*. New Delhi. Anmol Publication. 1993; **2** : 187-202.
19. Subba Rao, NV, Mitra SC. On land and freshwater molluscs of Pune District, Maharashtra. *Rec. Zool. Surv. India*. 1979; **75** : 1-37.
20. Subba Rao NV, Surya Rao KV, Manna RN. Wetland Ecosystem Series 1 : Molluscan Fauna of Chilka Lake. *Zool. Surv. India*. 1995; 391-468
21. Subharao NV. Fresh water molluscs of India. *Handbook Zoological Survey of India, Calcutta. XXIII* 1989; 1-289
22. Surya Rao KV, Mitra SC, Maitra S, Biswas. Tulika Molluscan fauna of Delhi, *State Fauna, Series 6*. *Zool. Surv. India*. 1997; 109-134.
23. Surya Rao KV, Mitra SC, Ramakrishna. Molluscan fauna of Kabar lake. *Wetland Ecosystem, Series 4*. *Zool. Surv. India*. 2002; 57-63